

Zebra/Quagga Mussel Surface Survey Protocol*

California Department of Fish and Game

*This protocol was adapted from the California Department of Water Resources *Zebra/Quagga Mussel Surface Survey Protocol*.

Description of Zebra and Quagga Mussels

The zebra mussel, *Dreissena polymorpha*, and the quagga mussel, *Dreissena bugensis*, are small mussels found only in freshwater. They look very similar to each other. They commonly have alternating light and dark brown stripes, but can also be solid light brown or dark brown. They have 2 smooth shells that are shaped a little bit like the letter “D”. These mussels are usually less than 2 inches in length. In new populations, most mussels are young and therefore very small (under ¼ inch long).

Quagga Mussel <i>Dreissena bugensis</i>	Zebra Mussel <i>Dreissena polymorpha</i>
	
<ul style="list-style-type: none"> • Shell: D-shaped and triangular; thin, fragile; smooth or shallowly ridged; solid light to dark brown or dark concentric rings; paler near hinge • Attaches to hard and soft surfaces 	<ul style="list-style-type: none"> • Shell: D-shaped and triangular; thin, fragile; smooth or shallowly ridged; solid light to dark brown or striped • Attaches to hard surfaces



Color variation in zebra and quagga mussels

Quagga and zebra mussels are freshwater mussels that can physically attach onto hard substrates. Like the mussels found clinging to the rocks along the California coastline, zebra and quagga mussels attach onto hard surfaces (e.g. pipes, screens, rock, logs, boats, etc.). They form colonies made up of many individuals attached onto an object and even onto each other. Small newly settled mussels feel like gritty sandpaper when attached to a smooth surface. Larger mussels will feel coarser (like a small pebble or sunflower seed) or be visually apparent.

Other Organisms Mistaken for Zebra/Quagga Mussels

Asian clam, Corbicula fluminea

People often mistake the very common Asian clam (also introduced) for zebra or quagga mussels. The Asian clam is widespread and abundant in California. It is brown and has ridges in concentric rings on its shells. The shells of older clams or of dead clams are white at the hinge (where the two shells join together). These clams do not attach onto surfaces. They live in mud or sand.



Snails and Freshwater Limpets

Small snails and freshwater limpets cling to hard substrates and can be mistaken for small juvenile mussels. They are similar in color and size to small zebra and quagga mussels. Snails have a spiral shape. Limpets have one shell and are flat. Zebra and quagga mussels attach on the edge of their shell and stick up and away from the surface.



Visual and Tactile Search for Zebra and Quagga Mussels

Gently run fingers over smooth surfaces, checking for gritty feeling or small “seed-like” or “pebble-like” objects. Areas likely to harbor mussels, if they are present, include:

- Dock floatation, buoys, mooring line, cables, rocks, concrete, logs/drift wood, vegetation, and anything that has been in the water for a long time.

- Pull up and inspect any substrate that is under water.
- Trap lines and any line or cable hanging in water.

Visually inspect all hard and soft substrates. Fan areas covered with silt to expose mussels.

Inspect dark areas (dark substrates and low light/shaded areas). Do not disturb private vessels or property.

Prime Areas to Search

Quagga and zebra mussels prefer dark substrates and low light/dark areas. They prefer concrete over other substrates. Search areas at or near boat ramps, gas dock, dock near marina store, other docks in high traffic areas, all concrete structures, and low flow areas.

Minimum Sample Size

The minimum number of linear feet to be searched per substrate is defined below. You can stop before meeting the minimum linear feet if quagga/zebra mussels are found in 3 or more locations within the survey location, or if all available substrate has been searched.

- Boat ramp bottom – 100ft if the ramp is at a marina, 200ft if the ramp is the only structure at the survey location.
- Shoreline - 100ft if at a marina, 200ft if at a survey location with only a boat ramp
- Dock - 200ft
- Mooring/dock lines (portion hanging in water) - 200ft
- Anchor/dock cable or chain (portion under water) - 100ft
- Concrete structures - 100ft
- Logs and woody debris – 100ft
- All accessible buoys

Make a notation in “Comments” section if minimum sample size requirements could not be met.

If Mussels are Found

Record the lat/long (in decimal degrees and use WSG 84) of the mussels’ location(s) and mark/describe location(s) on the back of the datasheet. Record the type of substrate(s) the mussel(s) was found on (for example, concrete, plastic, rope, chain, buoy, etc).

Make counts of mussels at up to 3 locations within the survey site. If more locations are found, make a note in the “Comments” section.

At each of the 3 mussel locations, take density estimates using one or both methods:

- Petri dish: place Petri dish over surface. Count all mussels within circle.
- Ruler: place ruler adjacent to mussels. Count all mussels within one inch of ruler.
- If you cannot see the mussels, count the mussels using touch. If entire ruler cannot be placed on surface, record length of ruler used.
- Collect 5 density estimates per mussel location.

Collect specimens (4-5). Place in Ziploc bag with label. Label should include location, lat/long, date, and name of collector. Seal and keep dry or put in freezer. Do not put water in the bag.

If other species of clams or mussels are found, collect specimens (1-2) and place in bag with collection label. Seal and keep dry or put in freezer. Do not put water in the bag.

Data Recording and Reporting

Datasheets are available at:

<http://www.des.water.ca.gov/docs/datasheet%20-%20surface%20survey.pdf>

If mussels are found, immediately contact the appropriate DFG regional mussel contact.

Every time a survey is made the data must be recorded on a datasheet before leaving the field. Absence information is as important to document as presence, so complete and submit a datasheet even if no mussels were found. Send datasheets to the appropriate DFG regional contact. All data will be entered into a data reporting system and the datasheets will be retained on-site.

DFG Regional Office Contacts for Quagga Mussel Monitoring

Region 1 – Northern Region

Counties: Del Norte, Humboldt, Lassen, Mendocino, Modoc, Shasta, Siskiyou, Tehama, and Trinity
601 Locust Street, Redding, CA 96001
L. Breck McAlexander
LMCALEXANDER@dfg.ca.gov
Office: (530) 225-2317
Fax: (530) 225-2381

Region 2 – North Central Region

Counties: Alpine, Amador, Butte, Calaveras, Colusa, El Dorado, Glenn, Lake, Nevada, Placer, Plumas, Sacramento, San Joaquin, Sierra, Sutter, Yolo and Yuba
1701 Nimbus Road, Rancho Cordova, CA 95670
Jason Roberts
JDROBERTS@dfg.ca.gov
Office: (916) 358-2895
Fax: (916) 358-2912

Region 3 – Bay Delta Region

Counties: Alameda, Contra Costa, Marin, Napa, Sacramento, San Mateo, Santa Clara, Santa Cruz, San Francisco, San Joaquin, Solano, Sonoma, and Yolo
4001 North Wilson Way, Stockton, CA 95205
Catherine Mandella
CMANDELLA@dfg.ca.gov
Office: (209) 942-6107
Fax: (209) 946-6355

Region 4 – Central Region

Counties: Fresno, Kern, Kings, Madera, Mariposa, Merced, Monterey, San Benito, San Luis Obispo, Stanislaus, Tulare and Tuolumne
1234 E. Shaw Avenue, Fresno, CA 93710
Mark Watson
MWATSON@dfg.ca.gov
Office: (559) 243-4017 x244
Fax: (559) 243-4022

Region 5 – South Coast Region

Counties: Los Angeles, Orange, San Diego, Santa Barbara and Ventura
4949 Viewridge Avenue, San Diego, CA 92123
Daniel Schrimsher
DSCHRIMSHER@dfg.ca.gov
Office: (858) 467-6926
Fax: (858) 467-4299

Region 6 – Inland Deserts Region

Counties: Imperial, Inyo, Mono, Riverside and San Bernardino
P.O. Box 2160, Blythe, CA 92226
David Vigil
DVIGIL@dfg.ca.gov
Office: (760) 922-4928
Fax: (760) 922-5638
Rick Francis
RFRANCIS@dfg.ca.gov
Cell: (760) 828-7582
Fax: (760) 922-5638